Fractional Cyclical Structures and Business Cycles in the Specification of the US Real Output

by L.A. Gil-Alana University of Navarra, Pamplona, Spain

Abstract

The issue in this paper is to analyse the business cycle frequencies in the US real output. However, instead of using classical approaches based on linear and non-linear models, we use a specification of fractional cyclical integration, which is based on Gegenbauer processes. We apply a procedure that permits us to test roots with integer and fractional orders of integration at fixed frequencies over time and thus, it permits us to approximate the length of the cycles. The results, based on the first differenced data, show that the cycles have a duration of about four years and a half, with an order of integration higher than 0 but smaller than 0.5, being thus stationary but with a component of long memory behaviour. Comparing this model with those based on ARIMA (and ARFIMA) models, we show via simulations that the fractional cyclical structure can better describe the business cycle features of the data.

Keywords: Gegenbauer processes; Business cycles; Long memory; US output.

JEL classification: C22.

1. Introduction

The existence of cycles in macroeconomic time series is a well-known stylised fact. However, its appropriate way of modelling is a matter that still remains controversial. With the development of the National Bureau of Economic Research (NBER)'s project of "Measurement without Theory" and the first extensive study of Burns and Mitchell (1946) on the American Economy, business cycles and their features have constituted a direct object of empirical analysis. Numerous studies have tried to describe them and to consider their stability over time. Romer (1986, 1994), Diebold and Rudebush (1992) and Watson (1994) have, for example, explored data to know if fluctuations have been smoother (lower amplitude and longer duration) after the Second World War. Also, Neftci (1984), Hamilton (1989), Beaudry and

Koop (1993) investigated new business cycles features¹, showing that cycles exhibit an asymmetry in their phases: recessions being deeper and shorter than expansions. Other authors, such as Candelon and Henin (1995) characterised the distributions of these features via bootstrapped simulation of simple linear (ARIMA) models for GDP. They could then locate the observed features of the last cycle and conclude that they are rather normal. A step further, Hess and Iwata (1997) used them as benchmarks to gauge the adequacy of macroeconomic stochastic time series models. They replicate via Monte-Carlo simulations different models for GDP. Then, they build for each model the distribution of the business cycle features and compare them to the historical business cycle characteristics. The best model is then selected as the one, which best replicates the historical features. Three types of linear models, namely, integrating a stochastic trend (ARIMA), a deterministic trend and a segmented trend (as in Perron, 1989) as well as several nonlinear ones (SETAR, Markov-Switching and Beaudry and Koop, 1993) are considered. They conclude that complex non-linear or linear models do not better replicate business cycle features than a simple linear ARIMA(1,1,0). Such a conclusion appears to be rather destructive for recent attempts, which have tried to better model GDP. Pesaran and Potter (1997) consider a nonlinear time series model to examine the non-linearity in US output, while Candelon and Gil-Alana (2004) use fractionally ARIMA (ARFIMA) models to characterise the business cycle components of the US and the UK GDP.

Another issue is to estimate the length of the cycles. In modern business cycle research, both empirical and theoretical macroeconomic and modelgenerated time series are detrended before analysis of the business cycle component. A vast majority of researchers use the Hodrick-Prescott (1997) filter (HP-filter) or Baxter and King's (1999) band-pass filter, and most authors conclude that business cycles have a duration of about six years. The HP filter has been interpreted as an approximation to an ideal high pass filter, eliminating frequencies of 32 quarters or greater. (See, e.g., Prescott, 1986, and King and Rebelo, 1993). Researchers relying on other methods often share this view about the duration of the business cycle component. Baxter and King (1999) construct a band-pass filter designed to extract cycles with duration between 1.5 and 8 years. Englund et al. (1992) and Hassler et al. (1994) use a band-pass filter in the frequency domain to extract cycles with duration between 3 and 8 years. Similar conclusions are obtained in Canova (1998), Burnside (1998), King and Rebelo (1999) and others.

In this article, we model the business cycle characteristics of the US real GDP from a different time series perspective, and consider the cycle as an

¹ These features integrate the third moment of the cycle as well as the conditional asymmetry in mean.

additional component to the trend and to the seasonal structure of the series. Similarly to these other two components, deterministic cycles (based in this context, on trigonometric functions of time) were shown to be inappropriate for modelling most macroeconomic time series. Harvey (1985) and others proposed stochastic cycles, and Gray et al. (1989, 1994) generalised them to allow for long memory. In particular, these authors considered processes like:

$$(1 - 2\mu L + L^2)^d x_t = u_t, \qquad t = 1, 2, ...,$$
(1)

where L is the lag-operator (Lx_t = x_{t-1}), d can be any real number, and where u_t is an I(0) process, defined, in the context of the present paper, as a covariance stationary process with spectral density function, which is bounded and bounded away from zero at any frequency on the spectrum. Clearly, when d = 0 in (1), x_t = u_t, and a "weakly autocorrelated" x_t is allowed for, as opposed to the case of d > 0 when the process is said to be "strongly autocorrelated" or also called "strongly dependent", so-named because of the strong association (in the cyclical part) between observations widely separated in time. Gray et al. (1989) showed that x_t in (1) is stationary if $|\mu| < 1$ and d < 0.50 or if $|\mu| = 1$ and d < 0.25. They also showed that the polynomial in (1) can be expressed in terms of the Gegenbauer polynomial C_{i,d} (μ), such that for all d \neq 0,

$$(1 - 2\mu L + L^2)^{-d} = \sum_{j=0}^{\infty} C_{j,d}(\mu) L^j, \qquad (2)$$

where

$$C_{j,d}(\mu) = \sum_{k=0}^{[j/2]} \frac{(-1)^k (d)_{j-k} (2\mu)^{j-2k}}{k! (j-2k)!}; \qquad (d)_j = \frac{\Gamma(d+j)}{\Gamma(d)}$$

 $\Gamma(x)$ is the Gamma function, and a truncation will be required in (2) to make (1) operational. Thus, the process in (1) becomes:

$$x_{t} = \sum_{j=0}^{t-1} C_{j,d}(\mu) u_{t-j}, \quad t = 1, 2, \dots,$$
(3)

and when d = 1, we have

$$x_{t} = 2 \mu x_{t-1} - x_{t-2} + u_{t}, \quad t = 1, 2, ...,$$
(4)

which is a cyclical I(1) process with the periodicity determined by μ . Tests of (4) based on autoregressive (AR) alternatives were proposed amongst others by Ahtola and Tiao (1987). Their tests are embedded in an AR(2) process of form:

$$x_{t} = \phi_{1} x_{t-1} + \phi_{2} x_{t-2} + u_{t}, \qquad (5)$$

which, under the null hypothesis:

 $H_0: |\phi_1| < 2 \text{ and } \phi_2 = -1,$

(6)

becomes the cyclical I(1) model (4). Unit root cycles were also examined by Chan and Wei (1988), Chung (1996) and Gregoir (1999a,b), who derive the limiting distribution of least squares estimates of AR processes with complex-conjugate unit roots, with inference based on parametric estimates.

In this article we use the fractional structure (1) for testing cyclical roots with integer and fractional orders of integration in raw time series. The use of this parametric approach to investigate the long run behaviour of time series consists of testing a parametric model for the series and relying on the long run implications of the estimated model. The primary advantage is the precision gained by focusing on the information in the series through the parameter estimates. A drawback is that the parameter estimates are sensitive to the class of models considered and may be misleading because of misspecification. However, the possibility of misspecification with parametric models can never be settled conclusively, and the problem can be addressed by considering a large class of models. For this purpose, we employ a version of the tests of Robinson (1994) that permits us to test long memory cyclical models. An advantage of this procedure is that is based on the Lagrange Multiplier (LM) priniciple and therefore we do not have to estimate any long memory fractional (cyclical) parameter. These tests are briefly explained in Section 2. Section 3 applies them to the quarterly structure of the US real output. In Section 4, we make some simulations comparing the business cycle characteristics of several models, while Section 5 contains some concluding comments.

2. The testing procedure

Following Bhargava (1986), Schmidt and Phillips (1992) and others on parameterization of unit-root models, Robinson (1994) considers the regression model:

$$y_t = \beta' z_t + x_t \qquad t = 1, 2, ...,$$
 (7)

where y_t is a given raw time series; z_t is a (kx1) vector of exogenous variables; β is a (kx1) vector of unknown parameters; and the regression errors x_t are such that:

$$\rho(L; \theta) x_t = u_t \qquad t = 1, 2, ...,$$
(8)

where ρ is a given function, which depends on L and the (px1) parameter vector θ , adopting the form:

$$\rho(L;\theta) = (1-L)^{d_1+\theta_1} (1-L^4)^{d_2+\theta_2} \prod_{j=3}^p (1-2\cos w L + L^2)^{d_j+\theta_j}$$
(9)

for real given numbers $d_1, d_2, \dots d_p$, and integer p, and where u_t is an I(0) process². Under the null hypothesis:

 $H_o: \theta = 0,$

(10)

(9) becomes:

$$\rho(L; \theta = 0) = \rho(L) = (1 - L)^{d_1} (1 - L^4)^{d_2} \prod_{j=3}^{p} (1 - 2\cos w L + L^2)^{d_j}$$
(11)

This is a very general specification that permits us to consider different models under the null. For example, if $d_1 = 1$ and $d_j = 0$ for $j \ge 2$, we have the classical unit-root model (Dickey and Fuller, 1979; Phillips and Perron, 1988; or the alternative in Kwiatkowski et al., 1992, etc.) and, if d_1 is a real value, the fractional models examined in Diebold and Rudebusch (1989), Baillie (1996) and others. Similarly, imposing $d_2 = 1$ and $d_j = 0$ for $j \ne 2$, we have the seasonal unit root model (Dickey, Hasza and Fuller, 1984, Hylleberg et al., 1990, etc.) and if d_2 is real, the seasonal fractional model analysed in Porter-Hudak (1990) and Gil-Alana (2002). Finally, if $d_1 = d_2 = 0$ and $d_3 = 1$, we have the unit root cycles of Ahtola and Tiao (1987); if d_3 is real, the model in Gray et al. (1989) and, if j > 3, the k-factor Gegenbauer processes studied in Ferrara and Guégan (2001).

In this article, we are concerned with the cyclical structure of the series and thus, we take $d_j = 0$ for $j \neq 3$ and $d_3 = d$. In such a situation (9) becomes:

$$\rho(L;\theta) = (1 - 2\cos wL + L^2)^{d+\theta}$$
(12)

and similarly (11), $\rho(L) = (1-2\cos wL + L^2)^d$. Plugging (12) in (7) and (8), y_t follows, under the null, a cyclical I(d) model of the form advocated by Gray (1989, 1994), and imposing d = 1, y_t appears as the unit root cycles proposed by Ahtola and Tiao (1987) and others.

We next describe the test statistic. We observe $\{(y_t, z_t), t = 1, 2, ..., n\}$, and suppose that the I(0) u_t in (8) have spectral density given by:

$$f(\lambda;\tau) = \frac{\sigma^2}{2\pi} g(\lambda;\tau), \quad -\pi < \lambda \leq \pi,$$

where the scalar σ^2 is known and g is a function of known form, which depends on frequency λ and the unknown (qx1) vector τ . Based on H_o (10), the residuals in (7); (8) and (12) are

$$\hat{u}_t = (1 - 2\cos wL + L^2)^d y_t - \hat{\beta}' s_t, \qquad (13)$$

² In Robinson (1994), the function $\rho(L;\theta)$ is slightly different. However, we have preferred to use this specification to stress the different components of the series, i.e., the trend, the seasonal and the cycles.

where
$$\hat{\beta} = \left(\sum_{t=1}^{n} s_t s_t'\right)^{-1} \sum_{t=1}^{n} s_t (1 - 2\cos wL + L^2)^d y_t,$$

with $s_t = (1 - 2\cos wL + L^2)^d z_t.$

Unless g is a completely known function (e.g., $g \equiv 1$, as when u_t is white noise), we have to estimate the nuisance parameter τ , for example by $\hat{\tau} = \arg \min_{\tau \in T^*} \sigma^2(\tau)$, where T^* is a suitable subset of \mathbb{R}^q Euclidean space, and

$$\sigma^{2}(\tau) = \frac{2\pi}{n} \sum_{s=1}^{n-1} g(\lambda_{s};\tau)^{-1} I_{\hat{u}}(\lambda_{s}), \quad \text{with}$$
$$I_{\hat{u}}(\lambda_{s}) = \left| (2\pi n)^{-1/2} \sum_{t=1}^{n} \hat{u}_{t} e^{i\lambda_{s}t} \right|^{2}; \quad \lambda_{s} = \frac{2\pi s}{n}.$$

The test statistic, which is derived via Lagrange Multiplier (LM) principle, takes the form:

$$\hat{R} = \hat{r}^2;$$
 $\hat{r} = \left(\frac{T}{\hat{A}}\right)^{1/2} \frac{\hat{a}}{\hat{\sigma}^2},$ (14)

where

$$\hat{a} = \frac{-2\pi}{n} \sum_{s=1}^{*} \psi(\lambda_s) g(\lambda_s; \hat{\tau})^{-1} I_{\hat{u}}(\lambda_s); \qquad \hat{\sigma}^2(\tau) = \sigma^2(\hat{\tau});$$

$$\hat{A} = \frac{2}{n} \left(\sum_{s=1}^{*} \psi(\lambda_s)^2 - \sum_{s=1}^{*} \psi(\lambda_s) \hat{\varepsilon}(\lambda_s)' \left(\sum_{s=1}^{*} \hat{\varepsilon}(\lambda_s) \hat{\varepsilon}(\lambda_s)' \right)^{-1} \sum_{s=1}^{*} \hat{\varepsilon}(\lambda_s) \psi(\lambda_s) \right),$$

$$\psi(\lambda_s) = \log |2(\cos \lambda_s - \cos w)|;$$
 $\hat{\varepsilon}(\lambda_s) = \frac{\partial}{\partial \tau} \log g(\lambda_s; \hat{\tau}),$

and the sum over * in the above expressions refers to all the discrete frequencies λ_s except the one with a pole in $\psi(\lambda)$.

Robinson (1994) showed that under certain very mild regularity conditions³,

³ These conditions are very mild regarding technical assumptions that are satisfied by the model in (7) - (9).

 $\hat{R} \rightarrow_{d} \chi_{1}^{2}$, as $n \rightarrow \infty$, (15)

Thus, we are in a classical large sample testing situation by reasons described in Robinson (1994). Because \hat{R} involves a ratio of quadratic forms, its exact null distribution can be calculated under Gaussianity via Imhof's algorithm. However, a simple test is approximately valid under much wider distributional assumptions. Thus, an approximate one-sided 100a%- level test of H_o (10) against the alternative: H_a: $\theta > 0$ ($\theta < 0$) will reject H_o if $\hat{r} >$ z_{α} ($\hat{r} < -z_{\alpha}$), where the probability that a standard normal variate exceeds z_{α} is α . Furthermore, he shows that the above test is efficient in the Pitman sense, i.e., that against local alternatives of form: H_a : $\theta = \delta n^{-1/2}$, with $\delta \neq 0$, the limit distribution is normal with variance 1 and mean that cannot (when ut is Gaussian) be exceeded in absolute value by that of any rival regular statistic. This version of Robinson's (1994) tests was examined in Gil-Alana (2001), and its performance in the context of unit root cycles was compared with Ahtola and Tiao's (1987) tests, the results showing that Robinson's (1994) tests outperform Ahtola and Tiao (1987) in a number of cases. Other versions of his tests have been applied to raw time series in Gil-Alana and Robinson (1997, 2001), testing for I(d) processes with the roots occurring respectively at zero and the seasonal frequencies. However, testing fractional cyclical models with the tests of Robinson (1994), this is the first empirical application, and one by-product of this work is its emergence as a credible alternative to the usual ARIMA (ARFIMA) specifications, which have become conventional in parametric modelling of macroeconomic time series.

3. The cyclical structure of the US real output

The time series data analysed in this section correspond to the quarterly, seasonally adjusted, real GDP in the US from 1947q1 to 2000q2, obtained from the Reserve Federal Bank of St. Louis' database. Alternatively, we could have employed seasonally unadjusted data, the reason for using adjusted data being that this series has been examined by many authors (e.g., Campbell and Mankiw, 1987; Diebold and Rudebusch, 1989; Hauser et al., 1992; Sowell, 1992a, and Koop et al., 2002). However, in all these papers they just concentrate at the long run or zero frequency and do not pay any attention to the cyclical structure underlying the series.





Figure 1 contains plots of the original series and its first differences, along with their corresponding correlograms and periodograms. We see that the original series increases over the sample period, though we also observe some apparent cyclical component in its behaviour. The nonstationary component of the series is also substantiated by the correlogram (with values decaying very slowly) and the periodogram (with a large peak around the smallest frequency). Taking first differences, the series may have a stationary appearance, though we still observe in the correlogram significant values even at some lags relatively away from zero, along with some apparent cyclical oscillation, which might imply the existence of some cyclical structure in the first differenced data.

The analysis below was first carried out based on the original data and there was not found any single evidence of fractional cycles across the sample. This is not surprising if we look at the periodogram of the original data, observing a large peak around the smallest frequency, which may suggest that first differences might be more appropriate when conducting this type of analysis⁴. In that respect, the model presented below might be related to the unobservable component (UC) models, (see, e.g., Harvey and Jaeger, 1993), where the long run component is specified as a random walk (or more generally, I(1)) trend⁵.

We also conducted the analysis based on the log-transformed data. Note that the log-transformation renders numerous statistical and interpretation advantages. Thus, its first differences can be interpretable as the GDP growth and it removes the exponential trend in the original data. The results using the log-transformation were less persistent (with smaller orders of integration) than the ones presented below. However, we have preferred to work with the original data since it is still unclear how the order of integration is affected in the context of Gegenbauer processes by this type of transformation⁶.

Denoting the differenced series by y_t , we employ throughout model (7), (8) and (12), with $z_t = (1,t)$ ', $t \ge 1$, (0,0)' otherwise, i.e.,

⁴ In fact, several tests for unit roots and I(0) stationarity (Dickey and Fuller, 1979; Kwiatkowski et al., 1992; and Hobijn et al, 1998) were conducted on the original data and the evidence was in all cases in favour of unit roots.

⁵ Similarly, we can relate this model with the Beveridge and Nelson's (1981) decomposition, which suggests, that any time series, which exhibits some form of nonstationarity of the type economic time series display, can be decomposed in two parts, a stationary component (the cyclical part) and a nonstationary one (the permanent component).

⁶ Another argument that may be presented in favour of the log-transformed series is that whilst the first differences show volatility that is increasing over time, the log-differenced series shows a decline in volatility that has been subject in the recent empirical work (Kim and Nelson, 1999; McConnell and Perez-Quiros, 2000; Blanchard and Simon, 2001; Stock and Watson, 2002; etc.). However, the issue of volatility is out of the scope of the present work. Moreover, Robinson's (1994) procedure used in the paper is supposed to be valid in the context of non-Gaussian and heteroscedastic disturbances.

$$y_t = \beta_0 + \beta_1 t + x_t, \qquad t = 1, 2, ...$$
 (16)

$$(1 - 2\cos w_i L + L^2)^{d+\theta} x_t = u_t, \qquad t = 1, 2, ...,$$
(17)

testing H_o (10) for values d = 0, (0.10), 2, and w_j = $2\pi/j$, j = 2, ..., n/2. Thus, j will indicate the number of periods required to complete the whole cycle. In other words, we test the order of integration of the cyclical polynomial at each of the frequencies of the process. Initially, we assume that $\beta_0 = \beta_1 = 0$ a priori, i.e., we do not consider deterministic regressors in the undifferenced regression (16). Then, we also examine the cases of β_0 unknown and $\beta_1 = 0$ a priori, (i.e., with an intercept), and both β_0 and β_1 unknown (i.e., with an intercept and a linear time trend), and report the results based on white noise and autocorrelated disturbaces⁷.

The test statistic reported across Table 1 corresponds to the one-sided statistic \hat{r} given by (14), for the case of white noise disturbances. However, instead of reporting the values for all possible combinations of d and j, we just report across the table, those cases where H_o cannot be rejected at the 95% significance level. We observe that the only values of d where H_0 cannot be rejected correspond to d = 0.10 and 0.20, with j constrained between 7 and 24 periods. The results are very similar for the three cases of no regressors, an intercept, and a linear time trend, and the fact that H_o is rejected for all d, if j is smaller than 7 or higher than 24, along with all j, if d = 0 or $d \ge 1$ 0.30, suggests that the optimal local power properties of the tests, shown by Robinson (1994), may be supported by reasonable performance against nonlocal alternatives. Another remarkable feature observed across this table is that for a given frequency i, the value of the test statistic monotonically decreases as d increases. Thus, for example, if j = 7, the values of \hat{r} with d = 0.10 are -0.305, -0.448 and 1.137 respectively for the cases of no regressors, an intercept, and a linear trend, and testing Ho, for the same j, with d = 0.20, the values reduce to -2.097, -1.971 and -1.561. This monotonicity is a characteristic of any reasonable statistic, given correct specification and adequate sample size. Thus, if H_0 (10) is rejected for a given d (and j) against alternatives of form: $\theta < 0$, an even more significant result in this direction should be expected when a higher value of d is tested. However, monotonicity is by no means necessary evidence of correct specification.

⁷ The FORTRAN codes of the programs used in this application are available from the author upon request.

| TABLE 1 | | | | | | |
|---|------|--------------------|-------------------|---------------------|--|--|
| Non-rejection values of the null hypothesis at the 95% significance level, using white noise ut | | | | | | |
| j | d | With no regressors | With an intercept | With a linear trend | | |
| 7 | 0.10 | -0.305 | -0.448 | 1.137 | | |
| 7 | 0.20 | -2.097 | -1.971 | -1.561 | | |
| 8 | 0.10 | -0.413 | -0.443 | 0.912 | | |
| 8 | 0.20 | -2.113 | -1.879 | -1.471 | | |
| 9 | 0.10 | 1.351 | 1.260 | 0.584 | | |
| 9 | 0.20 | -0.265 | -0.285 | -1.155 | | |
| 10 | 0.10 | 0.733 | 0.741 | 0.145 | | |
| 10 | 0.20 | -0.516 | -0.477 | -1.242 | | |
| 11 | 0.10 | 0.327 | 0.345 | -0.195 | | |
| 11 | 0.20 | -0.780 | -0.727 | -1.438 | | |
| 12 | 0.10 | -0.0002 | 0.024 | -0.395 | | |
| 12 | 0.20 | -1.013 | -0.956 | -1.546 | | |
| 13 | 0.10 | -0.154 | -0.173 | -0.424 | | |
| 13 | 0.20 | -1.078 | -1.041 | -1.484 | | |
| 14 | 0.10 | -0.514 | -0.489 | -0.742 | | |
| 14 | 0.20 | -1.522 | -1.458 | -2.113 | | |
| 15 | 0.10 | -0.758 | 0.686 | -0.905 | | |
| 15 | 0.20 | -1.998 | -1.584 | -2.321 | | |
| 16 | 0.10 | -0.953 | -0.835 | -1.018 | | |
| 16 | 0.20 | -2.115 | -1.740 | -1.990 | | |
| 17 | 0.10 | -1.056 | -0.939 | -1.095 | | |
| 18 | 0.10 | -1.072 | -0.991 | -1.048 | | |
| 19 | 0.10 | -1.401 | -1.256 | -1.358 | | |
| 20 | 0.10 | -1.403 | -1.252 | -1.340 | | |
| 21 | 0.10 | -1.546 | -1.380 | -1.454 | | |
| 22 | 0.10 | -1.765 | -1.528 | -1.566 | | |
| 23 | 0.10 | -1.865 | -1.616 | -1.637 | | |
| 24 | 0.10 | -1.621 | -1.423 | -1.472 | | |

In bold, the non-rejection values of the null hypothesis at the 95% significance level.



In order to have a more precise view about the non-rejection values obtained across Table 1, we have performed again the tests of Robinson (1994), but this time, for a range of values of d = 0, (0.01), 2. Figure 2 displays the (d, j) combinations where H_o cannot be rejected at the 95% significance



level. We see that the values of j range between 7 and 32, while d is in all cases smaller than 0.5, implying stationarity with respect to the cyclical component. These results are also in line with the literature mentioned in Section 1 about the duration of business cycles, with the periodicity constrained between 2 and 8 years.

The significance of the above results, however, might be in large part due to the un-accounted for I(0) autocorrelation in u_t . Thus, we also fitted other models, taking into account a weakly autocorrelated structure on the disturbances. We use AR(1) and AR(2) processes for u_t and the results, for the case of an intercept, are displayed in Figure 3. The first thing we observe here is that there is a larger proportion of non-rejection values compared with Figure 2 (white noise u_t) and, though j is again constrained between 7 and 32 periods, the values of d widely range between 0 and 0.6.

The results presented so far are not much conclusive since we observe many non-rejections, depending on j, d, the inclusion of deterministic trends and of the ARMA components. However, it should be noted that though the tests are unable to reject the null when d = 0, for some values of j, these hy-



potheses are in all cases "less clearly non-rejected"⁸ than when d is positive, suggesting that a component of long memory is present in relation with the cyclical structure.

Next, we try to be more specific about which may be the best model specification for this series, and the first thing we do is to examine the deterministic trends. We look at the coefficients β_0 and β_1 in (16) and, though not reported in the paper, the results showed that the coefficients corresponding to the time trend were insignificantly different from 0, in all cases where H_o could not be rejected. Note that these coefficients are all based on the null differenced model (13) and thus, standard t-tests apply. However, the coefficient for the case of an intercept was significant in many cases. Similarly for the weak dependence structure, we examine the coefficients corresponding to the AR components. Figure 4 resumes the (d, j) combinations of Figure 3 where all AR coefficients were significantly different from 0. We observe

⁸ By 'less clearly non-rejected', we mean that the value of the test statistic is closer to the rejection critical values.

| TABLE 2 | | | | | | | | | |
|--|--|--------|--------|--------|-----|--|--|--|--|
| Selected models for the case of $AR(1)$ ut with an intercept | | | | | | | | | |
| j | j d \hat{r} β_0 τ_1 Diagnostics | | | | | | | | |
| 15 | 0.51 | -0.004 | 34.330 | -0.923 | AC | | | | |
| 16 | 0.37 | -0.011 | 34.475 | -0.917 | AC | | | | |
| 17 | 0.27 | -0.013 | 34.589 | -0.912 | ABC | | | | |
| 18 | 0.19 | 0.017 | 34.680 | -0.906 | ABC | | | | |
| 19 | 0.14 | 0.020 | 34.754 | -0.902 | AC | | | | |
| 20 | 0.12 | 0.014 | 34.816 | -0.897 | AC | | | | |

A B and C means respectively that the models pass the diagnostic tests of no serial correlation (A), homocedasticity (B) and functional form (C).

| TABLE 3 | | | | | | | | | |
|---------|--|--------|--------|--------|--------|----|--|--|--|
| | Selected models for the case of $AR(2)$ ut with an intercept | | | | | | | | |
| j | d \hat{r} β_0 τ_1 τ_2 Diagnostic | | | | | | | | |
| 15 | 0.04 | -0.058 | 34.404 | -0.571 | -0.296 | А | | | |
| 16 | 0.04 | -0.057 | 34.408 | -0.404 | -0.153 | AB | | | |
| 17 | 0.02 | -0.030 | 34.412 | -0.251 | -0.044 | AB | | | |
| 18 | 0.02 | 0.063 | 34.414 | -0.247 | -0.037 | А | | | |

A B and C means respectively that the models pass the diagnostic tests of no serial correlation (A), homocedasticity (B) and functional form (C).

that the number of cases considerably reduces, especially for the AR(2) case: in both cases, j is constrained between 15 and 22, with d ranging between 0 and 0.50 for AR(1) u_t and between 0 and 0.20 for AR(2) u_t .

The results presented in Figures 3 and 4 correspond to the case of an intercept and AR(1) and AR(2) u_t respectively. Next, in Table 2 and 3, we choose for each j, the value of d, which produces the lowest $|\hat{r}|$ across d. The intuition behind this is that, for each j, the model with the closest statistic to zero will also be the one with the closest residuals to white noise. The third column of the tables reports the values of the statistics. The next two columns (and three in case of AR(2) u_t) correspond to the estimates of the intercept and the AR coefficients, while the last column reports some diagnostic tests carried out on the residuals of these selected models. An interesting feature observed across these tables is the fact that as we increase j, the value of d decreases, implying some type of competition between the length of the cycles and the orders of integration. The estimates of β_0 , τ_1 and τ_2 are significant in all cases.

Finally, we examine each of these potential models by looking at several diagnostic tests carried out on the residuals. In particular, we perform tests of no serial correlation, homoscedasticity and functional form, using Microfit.



Note that under the null hypothesis, the d_o -cyclical differences produce standard I(0) regressions (see equation (13)) and thus, standard techniques can be applied. The results are given in the last column of the tables. We see that the only models passing all the diagnostic tests on the residuals are those corresponding to AR(1) ut with j = 17 and 18. In the former case, the value of d is 0.27 while in the latter d = 0.19. Figure 5 displays the residuals for the two cases and we see that both have the appearance of white noise. Thus, we can conclude the analysis of this section by saying that the first differences of the US real output may be well described in terms of a Gegenbauer process, with the cycles occurring approximately every four years and a half, and with orders of integration ranging between 0.19 and 0.27, i.e., being stationary but with a component of long memory behaviour.

4. A simulation study: Comparisons across models

The objective in this section is to compare the fractional cyclical models of Section 3 with other approaches based on ARIMA and ARFIMA models, in terms of their ability to reproduce the business cycle features. For this purpose, we need to describe a rule to date the business cycles and define their characteristics. Numerous methods have been proposed in the literature. They can be based on direct data analysis (Burns and Mitchell, 1946), on expert judgement (NBER) or rely on the most recent econometric methods (Hamilton, 1994)⁹. In this paper, we have decided to consider exclusively classical cycles (directly extracted from the data in levels) in order to avoid statistical problems caused by the extraction of the cyclical component (See Canova, 1994). Besides, we apply the most common rule to date classical business cycles. This method was used in Candelon and Gil-Alana (2004), and it is at the basis of the famous program developed by Bry and Boschan (1971). It defines the phases of the business cycles as follows:

a) $y_{t-2} > y_{t-1} > y_t < y_{t+1}$, then there is a trough at t, where y_t is, for example, the GDP in levels.

b) $y_{t-2} < y_{t-1} < y_t > y_{t+1}$, then there is a peak at t.

c) When several identical turning points are detected consecutively, we retain the optimal one (i.e., the highest peak and the deepest trough).

This rule is very intuitive because it simply considers that a turning point occurs after two consecutive periods of expansion and recession. Such a rule consists of detecting a change in the slope of the process: Conditions a) and b) can be rewritten as $\Delta_j \ y_{t,j} > 0$ and $\Delta_j' \ y_{t+j}' < 0$, with j = 2 and j' = 1 for a peak (respectively as $\Delta_j \ y_{t+j} < 0$ and $\Delta_j' \ y_{t+j}' > 0$ for a trough). Such a definition insures that phases of the cycles have a minimum duration of 2 quarters and the completed cycles a minimum length of one year. This definition also presents the advantage of inducing an asymmetry in the length of the cycle phase, a property that is confirmed by historical data (Moore and Zarnowitz, 1982). On the contrary, we can not expect to detect an asymmetry in the amplitude of the phases, as the conditions on the change in slope are symmetric for troughs and peaks.

This method has been criticised in recent years. For example, it could exhibit not only major but also minor cycles. McNees (1991) and Webb (1991) propose to solve this problem via an increase in the reference period (for example, a peak could be characterised by 3 consecutive periods of growth over a year period). Candelon and Hénin (1995) have also noticed that this method leads to slight differences with the algorithms based on the detection of local optimum in the cases of growth cycles¹⁰. However, integrating these extensions in our dating algorithm will not alter the links between the degree of fractional cyclical integration and the business cycle

⁹ See Pagan and Harding (1999) for an exhaustive survey of the procedures for determining turning points. Also, Harding and Pagan (2003).

¹⁰ A local optimum is not a turning point for our methodology if it is preceded and followed by only one quarter of increase or decrease in the activity.



characteristics. We thus make the choice of simplicity and keep rules a) - c) as our dating algorithm.

From this dating, we have built five indicators (see Figure 6): the number of peaks (which corresponds to the number of cycles, as we consider that a cycle begins with a trough), the length of the cycles (period running between two successive troughs), the length and the amplitude of an expansion (period running from a trough to a peak), and the length and the amplitude of a recession (period running from a peak to a trough).

We have performed this dating algorithm for the series starting in 1960 in order to have better comparisons with reference studies. It is referred to the National Bureau of Economic Research (NBER) business cycle dating. It turns out in Table 4 that the algorithm leads to a nearly identical¹¹ dating except for the cycle (80:3-81:1), which is considered as minor in the official

¹¹ As NBER dating is performed for monthly data and our for quarterly observations, sometimes our results differ from a quarter.

| TABLE 4 | | | | | | | |
|---------|--------------|------------------|-----|---------------------|--------|--|--|
| | Busir | ess cycle datati | ior | 1 | | | |
| Country | Our datation | | | Reference datation* | | | |
| Country | Peak | Trough | | Peak | Trough | | |
| | 73:2 | 70:1 | | 73:4 | 70:4 | | |
| UNITED | 80:1 | 75:1 | | 80:1 | 75:1 | | |
| STATES | 81:1 | 80:3 | | | | | |
| SIAIES | 90:2 | 82:1 | | 90:3 | 82:4 | | |
| | 92.4 | 91.1 | | | 91.1 | | |

* Reference datation corresponds to the NBER datation.

| TABLE 5 | | | | | | | |
|---|--------------------------|--------------------------|-----------------------------|-----------------------------|--|--|--|
| Business cycle characteristics of the US real GDP | | | | | | | |
| Number of peaks | Mean length of expansion | Mean length of recession | Mean amplitude of expansion | Mean amplitude of recession | | | |
| 5 | 21.25 (3.44) | 5.00 (1.28) | 0.16 (0.04) | 0.02 (0.01) | | | |

Standard errors in parenthesis.

dating. In view of this, the above algorithm can be employed in simulated models when describing the business cycle characteristics.

Table 5 gathers the business cycle characteristics of the US real GDP. We notice that five major cycles occurred during the last 40 years. It also turns out that the expansions are longer and deeper than recessions. This stylised fact is generally known for classical cycles.

Table 6 reproduces the business cycle features of various models, which have been tried when modelling the US output. We start with AR(p)IMA(q) models with 1 and 2 differences. Here, we choose an ARIMA(1, 1, 1) and an ARIMA(1, 2, 1) respectively, and they have been chosen according to the Bayesian Information Criterion (BIC), from a group of sixteen potential models (with p and $q \le 3$). Besides, we consider fractional degrees of integration and thus, we also take into account an ARFIMA(0, 1.36, 0), which has also been selected by BIC¹². and the ARFIMA(3, 0.41, 2) proposed by

¹² Here, we use the Sowell's (1992b) procedure of estimating by maximum likelihood ARFIMA(p,d,q) models with p and $q \le 3$. Some theoretical properties of these stochastic processes can be found in Brockwell and Davies (1991), Odaki (1993) and Beran (1994). Moreover, when using integer differentiation we observe that if d = 1, the AR(1) coefficient is very large and close to 1, and, imposing d = 2, there is some evidence of overdifferentiation in the MA polynomial, implying that fractional processes with d lying between 1 and 2 (as in Sowell, 1992) might be more appropriate. Using the log-transformed data the persistence was found to be slightly smaller, which is in line with the results reported in Morley, Nelson and Zivot (2003) for the GDP growth rates.

| TABLE 6 | | | | | | | |
|---|----|----------------------------------|--|---------------------------------------|-----------------------------|--|--|
| Simulated business cycle characteristics of several models for the U.S. real GDP | | | | | | | |
| Model: ARIMA(1, 1, 1), $\phi = 0.95; \ \theta = -0.63$ | | | | | | | |
| Aver. Number of Peaks | | Mean length of expansion | Mean length of recession | Mean amplitude of expansion | Mean amplitude of recession | | |
| 12 | | 12.141 | 8.071 | 0.031 | 0.004 | | |
| (0,045) | | (3.654) | (2.321) | (0.010) | (0.043) | | |
| | | Model: ARIM | $A(1, 2, 1), \phi = 0.3$ | $30; \theta = -0.98$ | | | |
| Aver. Number of Peaks | | Mean length of expansion | Mean length of recession | Mean amplitude of expansion | Mean amplitude of recession | | |
| 3 | | 27.423 | 16.9742 | .2502 | 0.1707 | | |
| (0.035) | | (4.215) | (3.875) | (0.056) | (0.051) | | |
| Model: ARFIMA(0, d, 0), d = 1.36 | | | | | | | |
| Aver. Number of Peaks | | Mean length of expansion | Mean length of recession | Mean amplitude of expansion | Mean amplitude of recession | | |
| 9 | | 13.6392 | 8.8280 | 0.0994 | 0.0237 | | |
| (0.0761) | | (2.7195) | (2.4950) | (0.0101) | (0.0089) | | |
| Model: ARFIMA | (3 | , d, 2), $d = 0.41$, $\phi_1 =$ | = -1.18; $\phi_2 = 0.93; \phi_2 = 0$ | $\phi_3 = -0.51; \ \theta_1 = -0.29;$ | $\theta_2 = 0.81$ (Sowell) | | |
| Aver. Number of Peaks | | Mean length of expansion | Mean length of recession | Mean amplitude of expansion | Mean amplitude of recession | | |
| 14 | | 10.008 | 6.143 | 0.123 | 0.023 | | |
| (0.040) | | (3.113) | (3.129) | (0.008) | (0.0001) | | |
| Model: (16) and (17) with AR(1) u _t . $\beta_0 = 34.684$; $\beta_1 = 0$; $\tau = -0.906$; $j = 17$; $d = 0.27$ | | | | | | | |
| Aver. Number of Peaks | | Mean length of expansion | Mean length of recession | Mean amplitude of expansion | Mean amplitude of recession | | |
| 5 | | 26.504 | 9.110 | 0.013 | 0.041 | | |
| (0.0564) | | (3.123) | (3.452) | (0.006) | (0.034) | | |
| Model: (16) and (17) with AR(1) u _t . $\beta_0 = 34.754$; $\beta_1 = 0$; $\tau = -0.902$; $j = 18$; $d = 0.19$ | | | | | | | |
| Aver. Number of Peaks | | Mean length of expansion | Mean length of recession | Mean amplitude of expansion | Mean amplitude of recession | | |
| 5 | | 27.060 | 9.007 | 0.017 | 0.034 | | |
| (0.0487) | | (3.190) | (2.543) | (0.007) | (0.027) | | |

Sowell (1992b). Finally, the two selected models of Table 2, based on fractional cyclical integration, are also considered.

For each of these selected models, we simulate 2,500 series and compute their business cycle features. Their empirical means and variances are displayed in Table 6. Starting with the average number of peaks, we observe that the only models obtaining the exact number of peaks (5) are those based on the fractional cyclical structure. Also, the mean lengths of expansions and recessions are closer to the real business cycle characteristics with the latter models, though we loss part of the asymmetry in its behaviour. In relation with the amplitudes, the closest values to the true ones are obtained with the I(d) model. As a conclusion, we can summarise the results obtained in this section by saying that the fractional cyclical models of Section 3 can be used as alternative credible ways when modelling real output, performing better than the classical ARIMA (and ARFIMA) models when reproducing the business cycle features of the data.

5. Concluding comments

Economic theory in whatever format it comes has build-in the idea that cycles are stationary (if not in a wide sense, at least in a weak sense). That is, although there may be spillover of cyclical movements into the medium-long run, these are very small and nonstationary behaviour, if it exists, is due to the trend of the series. On the other hand, an enormous body of literature has also documented that a wide variety of indicators are persistent _ where persistence is defined in many ways, but roughly speaking it means that the process has long memory. These two points are a datum and any empirical specification that contradicts these facts should be look with suspicion. In this article we present an empirical model that faces these two empirical facts.

We have examined the cyclical structure of the US real output by means of a procedure of Robinson (1994) that permits us to test unit and fractional root cycles in raw time series. This method is based on Gegenbauer processes and tests the degree of integration of the cyclical component at fixed frequencies over time, which permits us to approximate the length of the cycles. The tests have standard (normal) null and local limit distributions, which is another distinguishing feature of these tests compared with other procedures. Moreover, they do not require Gaussianity for the asymptotic distribution, with a moment condition only of order 2 required. The tests were performed based on the first differenced data and the results showed that fractional cycles may be an alternative plausible way of modelling this series, with the order of integration ranging between 0.19 and 0.27 and a periodicity constrained between 15 and 20 quarters of a year.

These results have strong implications in terms of economic policy and planning inference. Thus, shocks affecting the cyclical structure will be mean reverting (i.e. their effects disappearing in the long run), which is in contrast with the long run effects, where shocks will persist forever. Then, stronger policy actions will be required in relation with the long run effects in comparison with the cyclical ones, assuming, for example, that the trend and the cycle are driven by the same shock. It is important to note, however, that as it was concluded by the literature of the late eighties, trend-cycle decompositions are not unique and there is not clear matching between restrictions coming from economic theory and the restrictions needed to achieve statistical identification. In that respect, the analysis of the forecasting ability of the proposed models is an area that should also be explored.

An argument that can be employed against the fractional cyclical model is that, contrary to seasonal cycles, business cycles are typically weak and irregular and are spread evenly over a range of frequencies rather than peaked at a specific value. However, contrary to that argument, we can explain that, in spite of the fixed frequencies used in this specification, the flexibility can be achieved throughout the first differencing polynomial, the ARMA components and the error term¹³. In that respect, the results presented here are completely in line with the literature on business cycle duration that says that cycles take place with a periodicity constrained between 3 and 8 years. Comparing these specifications with the classical ARIMA (and AR-FIMA) models, the results show that the former models better replicate the business cycle features of the US output. In fact, Hess and Isawa (1997) showed that the ARIMA models did it better than other linear and non-linear approaches (Perron, 1993; SETAR, Markov-Switching and Beaudry and Koop's 1993 non-linearity). Here, we have shown that fractional cyclical models can do it even better.

It would also be worthwhile proceeding to get point estimates for the fractional differencing parameter in the context of cyclical models. Some attempts have been made by Arteche and Robinson (2000) and Arteche (2002). However, the goal of this paper is to show that the Gegenbauer processes can be credible alternatives when modelling macroeconomic time series and, in that respect, the results presented in this paper lead us to unambiguous results, with the periodicity constrained between 15 and 20 periods and the order of integration higher than 0 but smaller than 0.5, implying stationarity and long memory in relation with the cyclical component.

The article can be extended in several other directions. First, the key results in the paper are based on first differenced data, and though the unit root null hypothesis is not rejected in the original series, it might also be of interest to perform other versions of Robinson's (1994) tests, which are larger in terms of model specification, for example, testing simultaneously the orders of integration at zero and the cyclical frequencies. However, in doing so, we lost the comparisons with other previous works, which are all based on first differences. Moreover, the existence of a unit root implies a stochastic trend and thus, the model can be alternatively written in the form of an orthogonal trend-cycle decomposition with an ARMA cycle which does not exhibit long memory. This specification is not nested in the model presented here, but it might be an alternative way of modelling its behaviour. Another potential

¹³ Bierens (2001) also use a model of this sort with d = 1, to test for the presence of business cycles in the annual change of monthly unemployment in the UK.

drawback of the present work is that it is based on an univariate model, with the limitation that it imposes in terms of theorizing, policy-making or forecasting. Theoretical models and policy-making involve the relationships between many variables and forecast performance can be improved through the use of many variables (e.g., factor based forecasts based on data involving hundreds of time series beat univariate forecasts, as described, for example, in Stock and Watson, 2002). However, the univariate work has relevance in the context of business cycles, firstly because different time series may have different amplitudes and different orders of integration, and there is not yet theoretical econometric models that permit us to examine cyclical fractional models in a multivariate framework. In that respect, the present paper can be considered as a preliminary step in the analysis of business cycles from a different time series perspective. Finally, the issue of data mining is another worry for economists when looking at time series models. There are so many possible models that may be relevant and so many modelling choices that econometricians are almost sure to find something purely by data mining. For this reason, sequential testing and other procedures based on information criteria are widely distrusted, and model averaging methods have become very popular. Thus, it might also be worthwhile to broaden the class of models under consideration and address the data mining problem, along with other issues (e.g., structural breaks) using averaging approaches. With respect to the latter issue, several papers conducted on the US output find little evidence of structural break in output, but strong evidence of structural break in output volatility, with the estimated break date around 1983. Diebold and Inoue (2001) point out that the evidence for long memory (at least at the zero frequency) in series that exhibit structural breaks may be spurious. We also conducted the analysis including a shift (and a slope) dummy for the break in 1983, and the results did not substantially differ from those reported across the paper.

References

- Ahtola, Juha and George C. Tiao (1987) Distributions of least squares estimators of autoregressive parameters for a process with complex roots on the unit circle. *Journal of Time Series Analysis* 8, 1-14.
- Arteche, Josu (2002) Semiparametric robust tests on seasonal and cyclical long memory series. *Journal of Time Series Analysis* 23, 1-35.
- Arteche, Josu and Peter M. Robinson (2000) Semiparametric inference in seasonal and cyclical long memory processes. *Journal of Time Series Analysis* 21, 1-25.
- Baillie, Robert T. (1996) Long memory processes and fractional integration in econometrics. *Journal of Econometrics* 73, 5-59.

- Baxter, Marianne and Roger M. King (1999) Measuring business cycles: Approximate band-pass filters for economic time series. *The Review* of Economics and Statistics 81, 575-593.
- Beaudry Paul and Gary Koop (1993) Do recessions permanently change output?. *Journal of Monetary Economics* 31, 149-163.
- Beran, Jan. (1994) *Statistics for long memory processes*. New York, Chapman and Hall.
- Beveridge, S. and Charles A. Nelson (1981) A new approach to decomposition of economic time series into permanent and transitory components with a particular attention to the measurement of the business cycle. *Journal of Monetary Economics* 7, 151-174.
- Bhargava, Alok (1986) On the theory of testing unit roots in observed time series. *Review of Economic Studies* 53, 369-384.
- Blanchard, Olivier and John Simon (2001) The long and large decline in US output volatility. *Brooking Papers on Economic Activity* 2001:1.
- Bierens, Herman J. (2001) Complex unit roots and business cycles: Are they real?, *Econometric Theory* 17, 962-983.
- Brockwell, Peter J. and Richard A. Davies (1991) *Time series: Theory and methods*. New York, Springer-Verlag.
- Bry, Gerhard and Charlotte Boschan (1971) *Cyclical analysis of time series:* Selected procedures and computer programs. New-York, NBER.
- Burns, Arthur F. And Wesley Mitchel (1946) *Measuring Business Cycles*. NBER, New-York.
- Burnside, A. Craig (1988) Detrending and business cycle facts. A comment, Journal of Monetary Economics 41, 513-532.
- Campbell, John Y. and N. Gregory Mankiw (1987) Are output fluctuations transitory?. *Quarterly Journal of Economics* 102, 857-880.
- Candelon, Bertrand and Luis A. Gil-Alana (2003) Fractional integration and buysiness cycle features. *Empirical Economics* 29, 343-359.
- Candelon, Bertrand and Pierre Y. Hénin (1995) La récession des années quatre-vingt dix a-t-elle été exceptionnelle?. *Economie et Prévision* 120, 51-71.
- Canova, Fabio (1994) Detrending and turning points. *European Economic Review* 38, 614-623.
- Canova, Fabio (1998) Detrending and business cycle facts. Journal of Monetary Economics 41, 475- 512.
- Chan, Y.N.H. and C.Z. Wei (1988) Limiting distributions of least squares estimates of unstable autoregressive processes. *Annals of Statistics* 16, 367-401.
- Chung, Ching-Fan (1996) Estimating a generalised long memory process. Journal of Econometrics 73, 237-259.

- Dickey, David A. and W.A. Fuller (1979) Distribution of the estimators for autoregressive time series with a unit root. *Journal of the American Statistical Association* 74, 427-431.
- Dickey, David A., D.P. Hasza, and W.A. Fuller (1984) Testing for unit roots in seasonal time series. *Journal of the American Statistical Association* 79, 355-367.
- Diebold, Francis X. and and Inoue, A., (2001) Long memory and regime switching, *Journal of Econometrics* 105, 131-159.
- Diebold, Francis X. and Glenn D. Rudebusch (1989) Long memory and persistence in aggregate output. *Journal of Monetary Economics* 24, 189-209.
- Diebold, Francis X. and Glenn D. Rudebusch (1992) Have the postwar economic fluctuations been stabilized?. *American Economic Review* 82, 993-1005.
- Englund, Peter; Torsten Persson and Lars E.O. Svensson (1992) Swedish business cycles: 1861-1988. *Journal of Monetary Economics* 30, 343-371.
- Ferrara, Laurent and Dominique Guégan (2001) Forecasting with k-factor Gegenbauer processes: Theory and applications. *Journal of Forecasting* 20, 581-601.
- Gil-Alana, Luis A. (2001) Testing stochastic cycles in macroeconomic time series. *Journal of Time Series Analysis* 22, 411-430.
- Gil-Alana, Luis A. (2002) Seasonal long memory in the aggregate output. *Economics Letters* 74, 333-337.
- Gil-Alana, Luis A. and Peter M. Robinson (1997) Testing of unit roots and other nonstationary hypotheses in macroeconomic time series. *Journal of Econometrics* 80, 241-268.
- Gil-Alana, Luis A. and Peter M. Robinson (2001) Testing seasonal fractional integration in the UK and Japanese consumption and income. *Journal of Applied Econometrics* 16, 95-114.
- Gray, Henry L.; Nien Fan Yhang, Wayne A. Woodward (1989) On generalized fractional processes. *Journal of Time Series Analysis* 10, 233-257.
- Gray, Henry L., Nien Fan Yhang, and Wayne A. Woodward (1994) On generalized fractional processes. A correction. *Journal of Time Series Analysis* 15, 561-562.
- Gregoir, Stéphane (1999a) Multivariate time series with various hidden unit roots. Part I: Integral Operator Algebra and Representation Theory. *Econometric Theory* 15, 435-468.
- Gregoir, Stèphane (1999b) Multivariate time series with various hidden unit roots. Part II: Estimation and testing. *Econometric Theory* 15, 469-518.

- Hamilton, James D. (1989) A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica* 57, 357-384.
- Hamilton, James D. (1994) *Time Series Analysis*. Princeton University Press.
- Harding, Don and Adrian Pagan (2003) A comparison of two business cycle dating methods. *Journal of Economic Dynamics and Control* 27, 1695-1698.
- Harvey, Andrew (1985) Trends and cycles in macroeconomic time series. Journal of Business and Economics Statistics 3, 216-227.
- Harvey, Andrew and Albert Jaeger (1993) Detrending, stylised facts and business cycle. *Journal of Applied Econometrics* 8, 231-247.
- Hassler, John, Petter Lundvik, Torsten Persson and Paul Soderlind, Paul (1994) *The Swedish business cycle: Stylized facts over 130 years*. in: Bergstrom and Vredin eds. Measuring and Interpreting Business Cycles, Oxford.
- Hauser, Michael A., Benedikt Pötscher and Erhard Reschenhofer (1992) Measuring persistence in aggregate output: ARMA models, fractionally integrated ARMA models and non-parametric procedures. Manuscript.
- Hess, Gregory D. and Shigeru Iwata (1997) Measuring and comparing business cycle features. *Journal of Business Economics and Statistics* 15, 432-444.
- Hobijn, Bart, Phillip Hans Franses and Marius Ooms (1998) Generalizations of the KPSS-test for stationarity. Econometric Institute Report 9802/A, Econometric Institute, Erasmus University Rotterdam.
- Hodrick, Robert J. and Edward C. Prescott (1997) Postwar US business cycles. An empirical investigation. *Journal of Money, Credit and Banking* 29, 1-16.
- Hylleberg, Svend, Robert F. Engle, Clive W. J. Granger and B. S. Yoo (1990) Seasonal integration and cointegration. Journal of Econometrics 44, 215-238.
- King, Robert G. and Sergio T. Rebelo (1993) Low frequency filtering and real business cycles. *Journal of Economics Dynamics and Control* 17, 207-231..
- King, Robert G. and Sergio T. Rebelo (2000) *Resuscitating real business* cycles. in Woodford and Taylor eds. Handbook in Macroeconomics, Amsterdam, Elsevier Science, Vol 1B, Chapter 14, pp.927-1007.
- Kim Chang-Jin and Charles Nelson (2001) Business cycle turning points. A new coincident index, and test duration dependence based on a dynamic factor model with regime switching, *The Review of Economics* and Statistics 80, 188-201.

- Koop, Gary; Eduardo Ley Levy, Jacek Osiewalskiand and Mark F. J. Steel (in press) Bayesian analysis of long memory and persistence using ARFIMA models. *Journal of Econometrics*.
- Kwiatkowski, Denis, Peter C.B. Phillips, Peter Schmidt and Youngcheol Shin (1992) Testing the null hypothesis of stationarity against the alternative of a unit root. *Journal of Econometrics* 54, 159-178.
- McConnell, M.M. and G. Perez-Quiros (2000) Output fluctuations in the United States: What has changed since the early 1980s?, *The American Economic Review 90*, 1464-1476.
- McNess, Stephen (1991) Forecasting cyclical turning points. The lessons of the last three recessions in K. Lahiri and G.H. Moore editors, Leading Economic Indicators CUP, 151-168.
- Moore, Geoffrey and Victor Zarnowitz (1982) Sequential signals of recession and recovery. *Journal of Business* 55, 57-85.
- Morley, James C, Charles C. Nelson and Eric Zivot (2003). Why are the Beveridge-Nelson and the Unobserved-Components decomposition of GDP, *The Review of Economics and Statistics* 35, 235-243.
- Neftçi, Salib N. (1984) Are economic time series asymmetric over the business cycle?. *Journal of Political Economy* 92, 307-328.
- Odaki, M. (1993) On the invertibility of fractionally differenced ARIMA processes. *Biometrika* 80, 703-709.
- Pagan, Adrian R. and Don Harding (1999) Dissecting the cycle: A methodological investigation. Melbourne Institute Working Paper, University of Melbourne.
- Perron, Pierre (1989) The great crash, the oil hock and the unit root hypothesis. *Econometrica* 77, 1361-1401.
- Perron, Pierre (1993) Trends, unit roots and structural change in macroeconomic time series. Unpublished manuscript, University of Montreal, Montreal.
- Pesaran, M. Hashem and Simon M. Potter (1997) A floor and ceiling model of US output. *Journal of Economics Dynamics and Control* 21, 661-695.
- Porter-Hudak, Susan (1990) An application of the seasonal fractional differenced model to the monetary aggregates. *Journal of the American Statistical Association* 85, 338-344.
- Prescott, Edward C. (1986) *Theory ahead of business cycle measurement*. Carengie-Rochester Conference Series on Public Policy 25, 11-44.
- Robinson, Peter M. (1994) Efficient tests of nonstationary hypotheses. Journal of the American Statistical Association 89, 1420-1437.
- Romer, Christina D. (1986) Is stabilization of the post-war economy a figment of the data? *American Economic Review* 76, 314-334.

- Romer, Christina D. (1994) Remeasuring business cycles. *Journal of Economic History* 54, 573-609.
- Schmidt, Peter and Peter Phillips (1992) LM tests for a unit root in the presence of deterministic trends. Oxford Bulletin of Economics and Statistics 54, 257-287.
- Sowell, Fallaw (1992a) Modelling long run behaviour with the fractional ARIMA model. *Journal of Monetary Economics* 29, 277-302.
- Sowell, Fallaw (1992b). Maximum likelihood estimation of stationary univariate fractionally integrated time series models. *Journal of Econometrics* 53, 165-188.
- Stock, James, H. and Mark W. Watson (2002). Macroeconomic forecasting using diffusion indexes. *Journal of Business and Economic Statistics* 20, 147-162.
- Watson, Mark W. (1994) Business-cycle durations and postwar stabilization of the U.S. economy. *American Economic Review* 84, 24-46.
- Webb, R. (1991) On predicting stages of business cycles in : K. Lahiri and G. Moore, eds, Leading economic indicators: New approaches and forecasting record. Cambridge University Press, 129-140.